

LARYNGO-VAC®

Product description

Laryngo-Vac® is a monovalent modified-live virus vaccine for the immunization of chickens to help protect against infectious laryngotracheitis (ILT).



Composition

Modified-live infectious laryngotracheitis virus

Formulation

Lyophilized modified-live viruses

Target species

Chickens

Indications

Laryngo-Vac is recommended for administration to healthy chickens to help protect against infectious laryngotracheitis (ILT).

Vaccination program

- Vaccinate birds when they are 4 weeks of age; or, if conditions dictate, birds may be vaccinated as early as 10 days of age.
- Revaccinate replacement birds at 16 to 20 weeks of age.
- This vaccine is recommended for the immunization of healthy birds. If an outbreak of ILT occurs on your operation, immediately institute a program of vaccination of birds farthest from the infected and work toward the infected birds.
- Dosage: one dose per bird.

Method of administration

Eye drop (intraocular), drinking water or coarse spray:
See the product insert for detailed instructions for vaccine administration.

Withholding period

21 days before slaughter

Presentation

- 10 x 1,000 doses
- 10 x 10,000 doses

Storage

- Store between 2°C to 8°C (35°F to 46°F).

Precautions

- Rehydrate the vaccine outside the pens.
- Rehydrate one vial at a time and use entire contents immediately after rehydration. For eye drop administration, use only 1,000-dose bottles.
- Do not stretch the number of doses to a larger number of birds than recommended.
- Vaccinate all susceptible birds on the premises at the same time.
- Vaccinate healthy birds only.
- Take care not to spill or spatter the vaccine.
- Use entire contents of vial when first opened.
- Inactivate unused contents before disposal.
- As a precautionary measure, curb traffic for 10 to 14 days between premises where birds have been vaccinated and areas where there are nonvaccinated susceptible birds.
- A mild conjunctival reaction may occur approximately four days following vaccination — more often in dry, dusty houses than in well-managed quarters. Such a reaction will disappear in about three days.
- Contains gentamicin as a preservative.

Contraindications

None, when used according to instructions.

Warnings

The use of this vaccine is subject to state laws, wherever applicable. Use only on premises with a history of infectious laryngotracheitis

SIZE	STANDARD PACK
1,000 doses	10
10,000 doses	10

Contact your local Zoetis representative for more information.